RESEARCH ARTICLE

James G. Ditty · Richard F. Shaw · Joseph S. Cope

Distribution of carangid larvae (Teleostei: Carangidae) and concentrations of zooplankton in the northern Gulf of Mexico, with illustrations of early *Hemicaranx amblyrhynchus* and *Caranx* spp. larvae

Received: 5 May 2003 / Accepted: 5 April 2004 / Published online: 7 July 2004 © Springer-Verlag 2004

Abstract We examined 1,825 bongo-net samples collected during Southeast Area Monitoring and Assessment Program (SEAMAP) ichthyoplankton surveys of United States Gulf of Mexico waters (1982–1986) for carangid larvae. Objectives were to describe the distribution of carangid larvae and to examine distribution patterns relative to areas of higher zooplankton volumes in order to reveal areas that may be important nurseries. Samples contained about 29,200 carangid larvae from 13 species or species complexes in 11 genera. Chloroscombrus chrysurusand Decapterus punctatus accounted for 91.7% of all larvae. We found that the 'scads' (D.punctatus, Trachurus lathami, and Selar crumenophthalmus) utilize temporally and/or spatially distinct spawning strategies to reduce co-occurrence of larvae. Samples contained fewer larvae than expected of the amberjacks (Seriola spp.), Caranx crysos, and C. hippos/ latus given the abundance of adults in the survey area, possibly due to inadequate sampling at appropriate times and locations, gear avoidance, or gear bias. Zooplankton displacement volumes (ZDVs) differed among regions and seasons and were inversely related to surface salinity and station depth. Differences among years were not significant. ZDVs were consistently highest near the Mississippi River delta and along the western Louisiana/ eastern Texas shelf, and moderately high during summer

Communicated by P. W. Sammarco, Chauvin

J. G. Ditty (⊠)

Southeast Fisheries Science Center, Galveston Laboratory, Fishery Ecology Branch, Taxonomy and Ecology Laboratory, NOAA Fisheries, 4700 Avenue U, Galveston, TX 77551, USA E-mail: Jim.Ditty@NOAA.gov

R. F. Shaw

Coastal Fisheries Institute, Center for Coastal, Energy and Environmental Resources, Louisiana State University, Baton Rouge, LA 70803-7503, USA

J. S. Cope

Virginia Institute of Marine Science, Chesapeake Bay Hall, P.O. Box 1346, Gloucester Point, VA 23062, USA

and fall along the shelf break, with localized pockets of elevated volumes over the eastern Gulf shelf. We suggest that *Chloroscombrus chrysurus,D. punctatus, T. lathami*, and possibly *Oligoplites saurus, Hemicaranx amblyrhynchus* and *Caranx crysos* spawn in frontal areas and/or along other hydrographic features that promote higher productivity. We provide new illustrations and descriptive information for the larvae of *H. amblyrhynchus* and discuss characters that separate early larvae of several species of *Caranx*.

Introduction

Members of the family Carangidae represent about 5% of the world's annual marine finfish landings, with major fisheries concentrating primarily on Decapterus punctatus (round scad) and Trachurus lathami (rough scad) (Leak 1977). Although Nakamura (1980) reported 24 species of carangids in the northern Gulf of Mexico (Gulf), information on the distribution and abundance of most carangids is inadequate to determine the feasibility of exploitation or greater exploitation. Exploratory surveys of the Gulf report potentially commercial quantities of adult D. punctatus, T.lathami, Caranx crysos (blue runner), and *Chloroscombrus chrysurus*(Atlantic bumper), but only D. punctatus are presently exploited (Juhl 1966; Bullis and Carpenter 1968; Bullis and Thompson 1970; Klima 1971). The small but abundant C.chrysurus may serve as an important food for predatory game and commercially important fishes (Reintjes 1979) and has value for reduction to fishmeal and fish oil (Leak 1977). Other carangids are highly regarded as food (pompano, Trachinotus spp.), game fish (Elagatis bipinnulata, rainbow runner; Seriola spp., amberjack), or bait (Caranx crysos, D. punctatus). Further, the amberjacks (Seriola spp.) currently face over-exploitation because of a dramatic increase in landings and the demand for fresh fish (Richards and McGowan 1989).

The distribution of carangid larvae in the Gulf is poorly known, with past studies focusing primarily on select taxa (Montolio 1976; Lyons 1978) or surveys of limited geographic and temporal coverage (Aprieto 1974; Houde et al. 1979; Leak 1981; Flores-Coto and Sanchez-Ramirez 1989; Sanchez-Ramirez and Flores-Coto 1993; Sanchez-Velasco and Flores-Coto 1994). A better understanding of spawning areas and seasonal spawning migrations requires basic information on the distribution of larvae, which is suggestive of the proximity of adult spawning concentrations (Houde 1982). Given that some fisheries exploit only spawning aggregations (Elwertowski and Boely 1971), a better understanding of the spatial and temporal distributions of carangid larvae may hasten development of fisheries (Leak 1981), permit better planning and development of ichthyoplankton surveys for spawning biomass estimation, and promote new fishery regulations for proper management of both new and existing fisheries. Our objectives were to describe the distribution of carangid larvae from United States Gulf of Mexico waters (an area that approximates the United States Exclusive Economic Zone/Fishery Conservation Zone), and to examine the distribution patterns of carangid larvae relative to areas of higher zooplankton volumes in order to reveal areas that may be important nurseries.

Material and methods

We examined 1,825 bongo-net samples for carangid larvae collected during Southeast Area Monitoring and Assessment Program (SEAMAP) ichthyoplankton surveys of United States Gulf of Mexico waters (i.e., the northern Gulf) from 1982 through 1986 (Fig. 1). National Marine Fisheries Service (NMFS) vessels sampled waters more than 10 m deep in a systematic grid of stations at about 55-km intervals. Coastal waters less than 10 m deep were sampled by participating Gulf States based on their sampling grid design. Texas did not participate.

Cruise survey procedures and zooplankton 'wet' volume displacement methods followed those outlined by Smith and Richardson (1977). Briefly, tows were made with a 60-cm bongo net (0.333-mm mesh) hauledobliquely from within 5 m of the bottom or from a maximum depth of 200 m. A flowmeter mounted in the mouth of each net estimated the volume of water filtered. Ship speed was about 45 m/min and net retrieval was 20 m/min. Tows extended a minimum of 10 min in clear water or 5 min if waters were turbid and stations were less than 95 m deep. Time of the tow (day/night) depended on when the ship occupied a station. Differences between the number of bongo-net samples displaced to estimate zooplankton displacement volumes (ZDV) (n=1,664) and the number of collections examined for fish larvae (n=1,825) result from missing data. Zooplankton samples were strained by sieve to remove large coelenterates and non-planktonic organisms larger than 2.5 cm, displaced, and sorted for fish larvae. We standardized resulting ZDV to ml/100 m³and natural-log transformed values before statistical analyses. Standard SEAMAP sampling protocol fixed samples in 10% buffered formalin followed by a transfer to 70% ethyl alcohol after 48 h.


We based larval identifications on existing descriptive literature (i.e., Aprieto 1974; Laroche et. al. 1984), where possible, and our own descriptive work. We did not separate larvae smaller than 5 mm standard length (SL) of jack crevalle (Caranx hippos) from those of horse-eye jack (C. latus), or yellow jack [C. (Carangoides)bartholomaei] from those of bar jack [C. (Carangoides) ruber] because early larvae of these species have not been adequately described. Instead, we grouped these species into complexes (C. hippos/latus or C. bartholomaei/ruber). In addition, current descriptions of small Seriola and *Trachinotus* larvae are insufficient to permit reliable species separation and we discuss these at the generic level. We considered SL synonymous with notochord length in preflexion larvae and recorded all measurements as mm SL. We excluded carangids larger than 13.9 mm from analyses because bongo nets do not efficiently sample larger sizes.

Annual SEAMAP ichthyoplankton surveys did not sample all regions, months, and depth zones uniformly. Consequently, we do not compare inter-annual variability because of substantial differences in the temporal and spatial distributions of stations. Instead, we calculated a density (number of larvae/100 m³) for each station and an arithmetic mean density for each month and region-season combination. We summed densities across stations within each category (i.e., month, season, depth, region, etc.) and divided the sum by the total number of stations in that category. Month was summed across years because not all months and regions were uniformly sampled over the study period.

We divided the northern Gulf into three geographic regions and three depth zones to examine carangid and zooplankton distribution patterns. Station depth served as a surrogate for distance from shore and the declining productivity gradient from the more eutrophic coastal waters to the oligotrophic open gulf waters. Inner continental shelf waters were less than 40 m deep; outer shelf waters were from 40 m to 180 m deep; and oceanic waters were those more than 180 m deep (Fig. 1). Differences among water masses, intensity of upwelling, and proximity to sources of freshwater and nutrient input can affect distribution and abundance patterns of zooplankton and fish larvae (Houde and Chitty 1976). The 40 m isobath, however, divided the continental shelf into areas of comparable size, which was convenient for evaluating differences in abundance between the inner and outer continental shelf. Latitude 24° 30' N was the southern boundary of the survey area in the eastern Gulf and 26° 00' N the southern boundary in the central and western Gulf (Fig. 1).

Hydrographic data were collected at the sea surface primarily by bucket. Water temperature was taken with

Fig. 1 Stations sampled during Southeast Assessment and Monitoring Program bongo-net surveys of the northern Gulf of Mexico (1982 through 1986) and defined areas



a hand-held thermometer and a water sample was returned to the laboratory for salinity analysis. We weighted hydrographic data by the number of larvae collected at each station before calculating the median and mean for each parameter, a method that emphasized the distribution of larvae rather than equally weighting stations. We are aware, however, that surface water temperature and salinity may be poor correlates of larval distribution patterns for those species not concentrated near the surface (Houde et al. 1979). Consequently, we used a percent cumulative frequency of 75% to define the hydrographic conditions (water temperature, salinity, and station depth) most often associated with the occurrence of larvae.

We grouped 3-month intervals into seasons based on monthly average water temperatures in the survey area as follows: Spring (March–May), Summer (June–August), Fall (September–November), and Winter (December–February). Cruises during April and May sampled primarily oceanic waters beyond the shelf in conjunction with NMFS annual larval tuna surveys. Cruises during March and from June through November sampled waters primarily over or immediately adjacent to the continental shelf (Fig. 1). We excluded winter from analyses, except in compiling hydrographic information for *Trachurus lathami*, because sampling was restricted primarily to waters near the Mississippi River delta in December.

Distribution and abundance patterns of carangid larvae permit inferences about adult distributions, spawning areas, and spawning times (Houde 1982). Accordingly, we compared the frequency distribution of stations where larvae were collected (positive catch stations) among regions, seasons, depth zones, and diel periods, respectively, with a likelihood ratio chi-square $[G^2, \alpha = 0.05]$ (Daniel 1990)] and evaluated significant differences through contrasts. The original model included season, region, depth zone, and diel period as factors. We retested the model after excluding diel and all three-way interactions, which had a minimal effect on the model. We used a non-parametric Wilcoxon signedranks test to assess the significance of diel differences in density of larvae [$\alpha = 0.05$ (Daniel 1990)] and canonical correlation to assess the relationship between density of larvae, ZDV, and environmental variables. We considered relationships 'strong' if correlations were greater than 0.70, 'moderately strong' if greater than 0.60, 'significant' if greater than 0.50, and 'weak' if less than 0.50. We ran the canonical analyses on all stations sampled for the season(s) and region(s) of peak abundance, which was species-specific. We used ArcView GIS 3.1 software, a kriging function, linear semi-variogram, and the ordinary Gaussian method to generate ZDV plots. Cell size was 2 km and had a fixed radius. An average of the five nearest neighbors provided the cell mean. We used a nonparametric Friedman test (Daniel 1990) to examine differences in ZDV among seasons, regions, and depth zones. A Tukey's multiple range test established the significance of distribution patterns.

mean was calculated by summing individual station densities and dividing by the number of stations sampled. Month was summed across years (1982–1986). Values in square brackets are total stations sampled; numbers in parenthesis next to mean densities are number of stations where larvae were collected Table 1 Monthly mean density of carangid larvae (number of larvae/100 m³) for all stations sampled during SEAMAP bongo-net surveys of the northern Gulf of Mexico. Monthly

Taxa	Spring			Summer			Fall			
	March [144]	April [164]	May [291]	June [354]	July [139]	March [144] April [164] May [291] June [354] July [139] August [224]	September [175] October [116] November [110] December [108]	October [116]	November [110]	December [108]
Chloroscombrus chrysurus	ı	1	0.2 (5)	44.5 (145)	40.0 (60)	42.9 (88)	33.6 (94)	0.3 (12)	a(1)	
Decapterus punctatus	^a (2)		0.2 (16)	2.3 (66)	0.3 (19)	1.5 (17)	8.2 (47)	2.9 (33)	1.1 (21)	< 0.1 (4)
Trachurus lathami	1.3 (36)	< 0.1 (17)	a(6)		,					< 0.1 (4)
Selar crumenophthalmus				< 0.1 (14)	< 0.1 (9)	0.4 (34)	0.1 (14)	0.2 (15)	0.2 (17))
Caranx crysos	ı			0.4 (53)	1.5(30)	0.5 (64)	a(8)			1
C. hippos/latus	1	< 0.1 (13)	0.6 (62)	0.4 (48)	0.1(16)	a(4)	0.1 (16)	^a (2)	1	1
Hemicaranx amblyrhynchus	ı			0.2 (6)	0.1(2)	< 0.1 (2)			1	1
Seriola spp.	ı	a(8)	a(7)	a(4)		a(3)	^a (3)	a(3)	a(1)	1
Selene spp.	I	Ì		0.3 (41)	0.3 (23)	0.4 (38)	0.2 (16)	a(3)	· 1	I
Oligoplites saurus	ı	ı	1	< 0.1 (5)	0.2 (4)	< 0.1 (3)		1	ı	1

'Mean density less than 0.05/100 m³

Results

Species distributions

Collections contained 29,203 carangid larvae from 13 species or species complexes in 11 genera. Numerically, *Chloroscombrus chrysurus* accounted for 82.9% of all larvae collected, followed by *D. punctatus* (8.8%), *Caranx hippos/latus* (2.9%), and *C. crysos* (1.9%). Other species contributed only 997 larvae or 3.5% to the numerical total. Larvae of *Chloroscombrus chrysurus* (n = 24,218) were abundant and widely distributed from May through November, but were most frequently collected during summer (Tables 1, 2). *C. chrysurus* were more abundant west than east of the Mississippi River delta, and relatively uncommon south of Tampa Bay, Florida (Fig. 2). Larvae were collected primarily over the inner shelf at surface water temperatures of 27°C and higher (Table 3).

D. punctatus (n = 2,582) was the second most frequent and abundant taxon. Round scad larvae were collected primarily off Florida during fall (Fig. 2) at stations less than 70 m deep, and at water temperatures and salinities of 25°C and 33.7 ppt or higher (Tables 1, 2, 3). Although D.punctatus was caught during all months sam-

Table 2 Mean density of carangid larvae (number/ 100 m³) for the northern Gulf of Mexico by season and region. Density by season and region was calculated by summing densities across stations and dividing by the total number of stations sampled. Numbers in parentheses are stations where

larvae were collected and total

stations sampled


(Table 1), all south of 27° 00′ N (i.e., below Tampa Bay, Florida) in waters more than 1,000 m deep. Larvae of T. lathami (n = 240) were collected during December and from March through May, and those of Selar crumenophthalmus (n = 256) were captured from April through November, but most frequently from August through November (Table 1). Both T. lathami and S.crumenophthalmus were taken primarily at stations deeper than 40 m; however, T.lathami was collected in cooler waters (median 21.4°C) than S.crumenophthalmus (median 28.2°C; Table 3). Most positive catch stations for S.crumenophthalmus were west of the Mississippi River delta and along the outer shelf. Sampling was too spatially restricted from December through February to assess adequately the gulf-wide distribution of T. lathami during winter, when larvae were abundant near the Mississippi River delta. SEAMAP samples contained at least three species of


pled, only four stations in December contained larvae

SEAMAP samples contained at least three species of *Caranx* (Fig. 3). We separated most early *Caranx* larvae into species or species complexes based on differences in pigmentation and body shape. *Caranx crysos* larvae have a series of discrete melanophores ventrally along the visceral mass and a pronounced gap in dorsal pigmentation between the nape and developing soft dorsal fin origin (Fig. 3a). Early *C. hippos/latus* and *C. bartholomaei/ruber*

Region	Spring (March–May)	Summer (June–August)	Fall (September–November)	Regional mean
Chloroscombrus	chrysurus			
Eastern	0.0 (0/139)	3.5 (19/171)	5.6 (22/109)	2.9 (41/419)
Central	0.1 (5/383)	53.9 (154/299)	10.2 (49/209)	20.5 (208/891)
Western	$0.0 \ (0/77)$	57.6 (120/247)	38.2 (36/83)	42.8 (156/407)
Season mean	< 0.1 (5/599)	43.1 (293/717)	14.7 (107/401)	
Decapterus punc				
Eastern	0.7 (19/139)	6.3 (70/171)	14.9 (65/109)	6.7 (154/419)
Central	^a (8/383)	0.1 (10/299)	1.3 (35/209)	0.4 (53/891)
Western	$0.0 \ (0/77)$	0.2 (22/247)	^a (1/83)	0.1 (23/407)
Season mean	0.2 (27/599)	1.6 (102/717)	4.7 (101/401)	
Trachurus latha				
Eastern	< 0.1 (12/139)	0.0 (0/171)	0.0 (0/109)	^a (12/419)
Central	0.5 (43/383)	$0.0 \ (0/299)$	0.0 (0/209)	0.2 (43/891)
Western	< 0.1 (7/77)	$0.0 \ (0/247)$	0.0 (0/83)	^a (7/407)
Season mean	0.3 (62/599)	0.0 (0/717)	0.0 (0/401)	
Selar crumenoph				
Eastern	^a (2/139)	0.2 (9/171)	0.2 (15/109)	0.1 (26/419)
Central	^b (2/383)	0.1 (12/299)	0.2 (24/209)	< 0.1 (38/891)
Western	^a (5/77)	0.3 (36/247)	0.2 (7/83)	0.2 (48/407)
Season mean	^a (9/599)	0.2 (57/717)	0.2 (46/401)	
Caranx crysos				
Eastern	^a (2/139)	^a (7/171)	^a (4/109)	^a (13/419)
Central	^a (5/383)	0.3 (62/299)	^a (3/209)	0.1 (70/891)
Western	0.0 (0/77)	1.4 (78/247)	^a (1/83)	0.9 (79/407)
Season mean	^a (7/599)	0.6 (147/717)	^a (8/401)	
C. hippos/latus		0	0	
Eastern	0.2 (20/139)	^a (3/171)	^a (4/109)	< 0.1 (27/419)
Central	0.3 (31/383)	0.3 (27/299)	0.1 (12/209)	0.2 (70/891)
Western	0.7 (24/77)	0.3 (38/247)	^a (2/83)	0.3 (64/407)
Season mean	0.3 (75/599)	0.2 (68/717)	< 0.1 (18/401)	
Selene spp.	0.0 (0.14.50)	2 (1 (1 = 1)	haven	2 (= 1 + 4 + 0)
Eastern	0.0 (0/139)	a(4/171)	^b (1/109)	^a (5/419)
Central	0.0 (0/383)	0.3 (39/299)	< 0.1 (11/209)	0.1 (50/891)
Western	b(1/77)	0.6 (59/247)	0.3 (7/83)	0.4 (67/407)
Season mean	^b (1/599)	0.3 (102/717)	0.1 (19/401)	

^aMean density less than 0.05/100 m³ ^bMean density less than 0.005/100 m³








Fig. 2a–c Distribution of the larvae of three species of carangid collected during Southeast Area Monitoring and Assessment Program bongo-net surveys of the northern Gulf of Mexico (1982 through 1986). Plots are for the month or season of greatest abundance (i.e., August for *Chloroscombrus chrysurus*; fall for *Decapterus punctatus*; and August for *Caranx crysos*)

larvae have a nearly contiguous series of melanophores ventrally along the visceral mass and dorsally along the trunk (Fig. 3d, h). Early *C.bartholomaei/ruber* larvae, however, have a 'deflected' dorsolateral row of pigments that runs diagonally across the trunk and pigment ventrally along the caudal peduncle that *C.hippos/latus* lack (Fig. 3d, h). Early larvae of *C.bartholomaei/ruber* also have a series of trunk melanophores along the lateral midline that extends from just behind the head to the caudal peduncle. This same lateral midline series begins behind the anus in *C. hippos/latus* larvae (Fig. 3d, e, h). After notochord flexion, the body profile of *C. hippos/latus* and *C.bartholomaei/ruber* becomes deeper and more rounded than in *C. crysos* of comparable length (Fig. 3).

Larvae of *C. hippos/latus*(n=848) and *C. crysos* (n=558) had similar spatial, but different temporal distributions. While both taxa were more abundant west than east of the Mississippi River delta and over the outer continental shelf, *C. hippos/latus* was collected primarily during May and June at 33 ppt and higher, and *C. crysos* was collected primarily from June to August at less than 33 ppt (Tables 1, 3). Both *C. hippos/latus* and *C. crysos* exhibited diel catch differences that favored higher catches during the day (P=0.001 and P=0.002, respectively). SEAMAP samples contained only six larvae of *C. bartholomaei/ruber* (3.5–5.5 mm), all caught during May 1982 at three stations 3,148 m deep near 26° 00′ N, 87° 05′ W (Fig. 4).

The larvae of other carangid species were relatively uncommon. Selene spp. (n=329) larvae were most abundant during summer over the outer continental shelf and west of the Mississippi River delta (Tables 1, 2, 3). Early bluntnose jack larvae (Hemicaranx amblyrhynchus; n=64; 2.0–5.3 mm SL) were collected at only 10 of 717 stations sampled during summer. All positive catch stations were located along the Louisiana/ Texas inner shelf in waters less than 26 m deep (nearly all during June at a salinity of 32 ppt; Tables 1, 3; Fig. 4). We found that pigment on the roof of the mouth and the bilateral and dorsal midline rows of pigment separated H. amblyrhynchus larvae (Fig. 5) from those of other co-occurring species. Larvae of Oligoplites

Table 3 Environmental data for stations where larvae were collected during SEAMAP bongo-net surveys (1982–1986) of the northern Gulf of Mexico. Differences in the total number of stations sampled between parameters result from missing data. Data were combined across seasons, regions, depth zones, and years

Species	Number of positive stations	Median salinity (ppt)	Salinity range (ppt)	Number of positive stations	Median temperature (°C)	Temperature range (°C)	Number of positive stations	Median station depth (m)	Depth range (m)
Chloroscombrus chrysurus	384	29.8	10.4–36.6	393	29.0	22.7–33.0	405	21	4–651
Decapterus punctatus	203	35.2	17.2-37.1	223	27.9	19.0-33.3	230	40	8-3,347
Trachurus lathami	29	36.2	9.2 - 37.7	59	21.4	14.3-27.6	62	84	16-3,475
Selar crumenophthalmus	107	35.3	18.9-37.4	108	28.2	21.0-31.1	112	70	11-3,219
Caranx crysos	157	33.1	23.4-36.7	158	29.0	24.1 - 33.8	162	80	9-3,148
C. hippos/latus	136	35.8	13.4-38.1	158	27.4	22.7 - 32.2	161	395	9-3,413
Hemicaranx amblyrhynchus	10	30.9	24.8-33.0	9	28.6	27.0-29.4	10	12	8-26
Seriola spp.	22	36.1	18.7-37.8	28	26.9	21.3-29.6	29	208	18-3,203
Selene spp.	118	32.3	13.4-36.6	119	28.9	23.5-30.9	122	50	7-1,793
Oligoplites saurus	12	31.9	20.2–36.0	2	29.2	27.8–32.0	12	17	11–59

Fig. 3 Early larvae of *C. crysos* and two *Caranx* complexes collected during SEAMAP bongo-net surveys of the northern Gulf of Mexico from 1982 through 1986: *Caranx crysos*: a 3.4 mm, b 4.8 mm, c 6.8 mm; *Caranx hippos/latus*: d 3.6 mm, e 4.7 mm, f 7.1 mm, g 13.2 mm; *C. bartholomaei/ruber*: h 3.8 mm, i 5.5 mm standard length





Fig. 4 Distribution of some less abundant carangids collected during SEAMAP bongo-net surveys of the northern Gulf of Mexico from 1982 through 1986. The distribution of Oligoplites saurus overlaps that of Hemicaranx amblyrhynchus in the lighter gray area



saurus (leatherjackets; n = 54; 1.8–5.3 mm) were collected from June through August, generally shoreward of the 20 m isobath (Fig. 4). All *Seriola* spp. larvae

(n=48) smaller than 5.0 mm could not be identified to species because dorsal and anal ray development was incomplete. *Seriola* larvae were widely distributed from




Fig. 5a-d *H. amblyrhynchus* (top to bottom 3.1 mm dorsal and lateral views, 4.0 mm, and 5.3 mm SL)

April through October and were collected primarily at stations more than 40 m deep (Tables 1, 3; Fig. 4). Samples contained two larvae of pompano (*Trachinotus* spp.) and one African pompano (*Alectis ciliaris*). Both *Trachinotus* larvae (2.2–2.3 mm) were caught during April 1984 at stations more than 1,800 m deep (Fig. 4). The only African pompano larva (7.5 mm) was collected during June 1982 off South Florida at 36.3 ppt, 28.2°C, and in waters 33 m deep (Fig. 4).

Table 4 Mean wet zooplankton displacement volumes (ml/100³) for the northern Gulf of Mexico from SEAMAP bongonet surveys (1982 through 1986). Mean volume was calculated by summing individual station densities and dividing by the number of stations sampled per region or depth zone. Month was summed across years

ND No data

Zooplankton displacement volumes

Zooplankton displacement volumes (ZDVs) differed among regions, with distance from shore, and seasonally (Table 4). Overall, the central and western Gulf had significantly higher ZDVs than did the eastern Gulf (Table 5). ZDVs over the inner shelf (less than 40 m) tripled from winter to spring and declined thereafter, whereas volumes remained generally consistent throughout the year over the outer shelf, and in oceanic waters. Overall, ZDV was inversely related to surface salinity (canonical loading = 0.690) and station depth (canonical loading = 0.955), and positively related to surface water temperatures, but not significantly so. Differences among years were not significant.

ZDVs were consistently higher near the Mississippi River delta and along the western Louisiana/east Texas shelf, and moderately high along the shelf break during summer and fall, with localized pockets of elevated volumes over the eastern Gulf shelf (Fig. 6a–d). Similarly, ZDVs were moderately high over the shelf break during summer and fall, and also over the Florida continental shelf during fall. ZDVs were high near the Mississippi River Delta during winter (Fig. 6d).

The distribution of *D. punctatus* larvae over the continental shelf during fall and of *Chloroscombrus chrysurus* over the inner shelf during August (Fig. 2) was significantly correlated with higher ZDVs (Table 6). Despite the large number of plankton samples taken over the study period (more than 1,800 samples), *C.chrysurus* and *D. punctatus* were the only carangids abundant enough to statistically examine distribution patterns relative to areas of higher ZDVs. Nevertheless, locations along the inner shelf where larvae of *H.amblyrhynchus* and *O. saurus* were abundant during summer (Figs. 4, 6b), and near the Mississippi River delta where *Trachurus lathami* was abundant during winter and spring had higher coexisting ZDV values.

Discussion

C. chrysurus, one of the most abundant and widely distributed carangids in the Gulf of Mexico, spawn primarily during summer in waters less than 40 m

	Spring			Sumn	ner		Fall		
	March	April	May	June	July	August	September	October	November
Region									
Eastern	11.7	9.7	11.9	36.4	15.0	27.4	18.0	24.2	20.0
Central	46.0	6.6	41.6	68.0	85.3	25.3	30.0	38.2	35.0
Western	ND	9.3	7.2	53.9	32.2	28.0	30.1	27.0	24.0
Depth zone									
< 40 m	58.2	120.0	181.6	80.1	76.8	49.9	35.6	52.4	42.5
41–180 m	35.3	27.3	20.5	22.8	14.0	22.5	15.7	16.0	14.4
> 180 m	13.8	6.3	8.0	7.2	6.2	7.7	6.4	4.8	7.6
Total stations	137	163	269	306	136	223	152	108	107

Table 5 Spatial and temporal distribution of wet zooplankton displacement volumes (ml/ 100 m^3) for the northern Gulf of Mexico. Tukey's multiple range test established the significance of distribution patterns. Different letters within categories indicate significant differences at $\alpha = 0.05$

	Total stations	Mean volume (SD in parentheses)	Coefficient of variation	Tukey's multiple range test
Region				
Eastern	417	21 (31)	144%	C
Central	843	42 (64)	152%	A
Western	404	31 (44)	142%	В
Season		. ,		
Spring	590	26 (50)	192%	В
Summer	707	44 (63)	144%	A
Fall	367	29 (31)	108%	В
Depth				
< 40 m	634	67 (73)	92%	A
40–180 m	485	21 (19)	92%	В
> 180 m	545	7 (7)	97%	C

deep (Table 7). The fact that Atlantic bumper grow more quickly during summer (Leffler and Shaw 1992; Sanchez-Ramirez and Flores-Coto 1998) when water temperatures and ZDVs are traditionally high should not be surprising given that water temperature and food availability influences growth rates. The overlapping spatial and temporal distributions of higher quantities of Atlantic bumper and ZDVs suggest that the availability of food over the inner shelf during summer may have a positive affect on Atlantic bumper survival.

We found that the three species of 'scad' (Decapterus, Trachurus, and Selar) spawn primarily over the outer continental shelf, but utilize different spawning strategies to reduce co-occurrence of larvae. T. lathami are coolwater spawners (generally 22°C or lower) and are abundant in the eastern (Leak 1977), central (Ditty and Truesdale 1984), and southern Gulf of Mexico (Flores-Coto and Sanchez-Ramirez 1989) during winter/early spring when large schools of adults are present along the outer shelf (Leak 1977). In contrast, Decapterus and Selar spawn in warmer waters (generally above 24°C), with D. punctatus more abundant in the eastern Gulf and S. crumenophthalmus more abundant west of the Mississippi River delta. The fact that schools of adult S. crumenophthalmus (Reintjes 1979) co-occur along the outer shelf/shelf break in the western Gulf when larvae are more abundant is consistent with the possibility that this area and possibly the southern Gulf (Flores-Coto and Sanchez-Ramirez 1989) may be a spawning and nursery area for bigeye scad.

The existence of *D. punctatus* larvae south of 27° 00′ N during all months sampled in this study, when combined with similar findings year-round by Lyons (1978) and Leak (1981), is consistent with probable continual spawning and a southward offshore migration of adults during late fall—early winter (Klima 1971; Hales 1987), followed by a northward onshore return during spring. This north—south migration suggests unsuitable spawning conditions above 27° 00′ N during winter and may account for a pattern that has been interpreted as bimodal spawning in *D. punctatus*. Alternately, spawning could be continuous but appear bimodal (spring and fall) if egg and larval mortality is

higher during the middle of the spawning season (McBride et al. 2002).

The temporal distribution patterns we found for *Caranx crysos* larvae differed from those of Montolio (1976), who identified large numbers of *C. crysos* in Gulf oceanic waters during April–May, with a secondary peak during August–September. Although SEAMAP bongonet surveys from 1982 through 1986 collected 455 samples in oceanic waters during April and May, only nine samples contained *C. crysos* larvae. We suspect that some, if not most, of Montolio's larvae smaller than 5 mm are *C.hippos/latus* or possibly *C.bartholomaei/ruber* based on the distribution of larvae of these two species complexes in this study; the description by McKenney et al. (1958) of *C. crysos* larvae; and pigmentation differences between Figs. 1 and 2 in Montolio (1976).

We were unable to separate confidently *C. bartholomaei* from *C.ruber* because collections contained only six larvae, all smaller than those described previously (i.e., *C. bartholomaei* larger than 6.0 mm, *C.ruber* larger than 12.4 mm; Berry 1959). Gill raker counts along the lower limb separate juveniles and adults of *C. hippos* (22–27) from *C. latus* (16–18), and *C. bartholomaei* (18–21) from *C.ruber* (31–35) (Berry 1959). Adult *C. bartholomaei* and *C. ruber* are rare in the survey area and spawn during spring and summer in oceanic waters (Berry 1959; Nakamura 1980).

Four species of *Seriola* (*S.dumerili*, *S. fasciata*, *S. rivoliana*, and *S. zonata*) occur in the Gulf, but larvae of only the banded rudderfish (*S. zonata*) and greater amberjack (*S. dumerili*) have been described (Aprieto 1974; Masuma et al. 1990; Tachihara et al. 1993). Amberjack larvae smaller than 5 mm were taken from April through November (Table 1), although some species may spawn year-round off south Florida (Leak 1981). Reproductive data suggest that *S. dumerili* spawn during late spring/early summer (Thompson et al. 1998) and that *S.fasciata* spawn during late summer/fall. The center of *S. fasciata*distribution may be off Louisiana (Thompson and Brown 1994). Reproductive data are not available for *S.rivoliana* and *S. zonata*.

Lookdown larvae (*Selene setapinnis*) are found primarily in the western region of the Gulf during summer/

Fig. 6a–d Wet zooplankton displacement volumes (in ml/100 m³) for the northern Gulf of Mexico from SEAMAP bongo-net surveys (1982 through 1986). Seasons were spring (March–May), summer (June–August), fall (September–November), and winter (December–February). Border delineates area sampled. Codes (ml/100 m³) are 1 (<25), 2 (25-49), 3 (50-74), 4 (75-99), 5 (100-150), 6 (>150)




Table 6 Distribution of the larvae of two species of Carangidae from the northern Gulf of Mexico (1982–1986) relative to concentrations of zooplankton and hydrographic conditions. All stations sampled within the defined region, season, and depth zone

were used in the canonical correlation analysis. Numbers in bold signify significant correlations. Correlations were 'strong' if greater than 0.70, 'moderately strong' if greater than 0.60, 'significant' if greater than 0.50, and 'weak' if less than 0.50

Taxon	Total stations sampled	$\frac{\text{Canor}}{R}$	$\frac{\text{nical}}{R^2}$	P	Region	Season	Depth zone	Left Root	Factor structure	Right Root	Factor structure
Decapterus punctatus	109	0.564	0.318	< 0.0001	East	Fall	<180 m	Density	-1.000	Zooplankton Temperature Salinity Station depth	- 0.797 -0.315 0.110 0.557
Chloroscombrus chrysurus	300	0.629	0.395	< 0.0001	Central and west	Summer	<40 m	Density	-1.000	Zooplankton Temperature Salinity Station depth	-0.790 -0.495 0.136 0.751

Table 7 Spatial and temporal distribution of the larvae of carangids known to occur in the northern Gulf of Mexico

Taxon	Adults ^a	Spawni	ing season	ı ^b		Regi	on		Depth	zone ^c		Literature cited ^d
		Spring	Summer	Fall	Winter	East	Central	West	Inner	Outer	Oceanic	
Chloroscombrus chrysurus Decapterus punctatus	C C	?	X X	X		X X	X	X	X X	X		1, 2, 3, 4, 5, 6 1, 2, 4, 5, 7, 8, 9
Selar crumenophthalmus Trachurus lathami	C C		X	X	X	X	X	X X		X X		1, 2, 4, 13, 19
Caranx hippos/latus	C C	X	X		Λ	Λ	X X	X		X X X	X	2, 3, 4, 5, 7, 10, 11, 12 1, 4
C. crysos C. bartholomaei/ruber	R		X X					X			X	1, 4, 5, 7, 14, 17 1, 18
Seriola dumerili S. fasciata	C C	X	X X	X		?	X X	?		?	?	20 21
S. rivoliana S. zonata	C C	X X	?	X		? X	X	?		? X	?	1 13
Hemicaranx amblyrhynchus Selene setapinnis			X X	X				X X	X	X		1, 15 1, 4
S. vomer	C	v	X X	X		?		X	v	X		1, 2, 4, 13
Oligoplites saurus Trachinotus carolinus	C C	X ?	Λ	X	X				X			1, 2, 4, 13, 16, 18 1, 16, 18
T. falcatus T. goodei	C R	?		X	X							1, 16, 18 18
Alectis ciliaris Elagatis bipinnulata Naucrates ductor Uraspis secunda	R C R	X ? ?	X ?	?							X	1, 2, 10, 18 13, 18 18, 22 18

^aC common, R rare, ? no information

early fall (Table 7), which is consistent with the higher reported abundance of lookdown further south off Mexico (Flores-Coto and Sanchez-Ramirez 1989; Flores-Coto et al. 2000). Although larvae of *Selene* were not particularly abundant in SEAMAP samples, larvae of the Atlantic moonfish (*S. vomer*) are common off south Florida (Aprieto 1974; Leak 1981).

The fact that *H. amblyrhynchus* larvae have only been collected along the western Louisiana/eastern Texas inner shelf (Table 7) and off Mexico (Flores-Coto et al. 1998) is consistent with the suggestion that adult bluntnose jack are uncommon in the northern Gulf (Nakamura 1980; Flores-Coto et al. 2001). Although

1990; 6 Katsuragawa and Matsuura 1992; 7 Lyons 1978; 8 Hales 1987; 9 Montolio 1976; 10 Finucane et al. 1979a, b; 11 Ditty and Truesdale 1984; 12 Leak 1977; 13 Aprieto 1974; 14 Goodwin and Finucane 1985; 15 Hoese 1965; 16 Houde et al. 1979; 17 Klima 1971; 18 Nakamura 1980; 19 Reintjes 1979; 20 Thompson et al. 1998; 21 Thompson and Brown 1994; 22 Fahay 1975

adult *O. saurus* are common (Nakamura 1980), ichthyoplankton surveys have collected relatively few larvae. Based on our findings and other information, *O. saurus* spawn during summer over the inner shelf (Table 7).


Three species of pompano (*Trachinotus carolinus*, *T. falcatus*, and *T. goodei*) occur in the study area, although bongo-net samples contained only two *Trachinotus* larvae (probably *T. carolinus*), both from oceanic waters during April. The presence of small larvae during April is consistent with a late winter/early spring spawning period for *Trachinotus* as has been suggested for United States Atlantic coast waters (Fields 1962; Fahay 1975). A proposed late winter/early spring

^bSpring March-May, summer June-August, fall September-November, winter December-February

^cLess than 40 m, 40–180 m, greater than 180 m

^dLiterature cited: 1 This study; 2 Leak 1981; 3 Ditty et al. 1988; 4 Flores-Coto and Sanchez-Ramirez 1989; 5 Shaw and Drullinger

Fig. 7 Surface water temperatures in the northern Gulf of Mexico during summer (June through August) based on samples collected during SEAMAP surveys from 1982 through 1986



spawning period is also consistent with finding juvenile Florida pompano (*T.carolinus*) and permit (*T. falcatus*) relatively common during spring and summer in highenergy surf zones of the Gulf (Ruple 1984). Collection of a 5.2 mm *T.falcatus* larva during August (Flores-Coto et al. 2000) suggests summer spawning in this species. *T. goodei* (palometa) are uncommon in the northern Gulf (Nakamura 1980).

Bongo-net samples contained only one *A. ciliaris* (African pompano) larva and no larvae of *E. bipinnula-ta*(rainbow runner), *Uraspis secunda*(cottonmouth jack), or *Naucrates ductor*(pilotfish). *A. ciliaris* are apparently rare and larvae have been collected only near the southern limit of the survey area during summer and early fall (Table 7). *E. bipinnulata* spawn in oceanic waters (Aprieto 1974) and adults are common offshore, whereas adults of *U. secunda* and *N.ductor* are rare (Nakamura 1980). We found no documented record of either *U. secunda* or *N. ductor* larvae from the survey area. Fahay (1975), however, collected two young *N. ductor* smaller than 12 mm off the United States Atlantic coast during May, suggestive of early spring spawning.

We found the larvae of several species of carangids (Chloroscombrus chrysurus, Trachurus lathami,D. punctatus, Caranx crysos,O. saurus, and H. amblyrhynchus) in hydrographically dynamic areas where ZDVs are traditionally higher (Arnold 1958; Bogdanov et al. 1968; Khromov 1969; Austin and Jones 1974; Houde and Chitty 1976; Howey 1976; Dagg et al. 1987). ZDVs are consistently higher in areas influenced by riverine output as evidenced by higher volumes near and west of the Mississippi and Atchafalaya River deltas (Fig. 6), a relationship consistent with fish adapting their spawning strategy to local oceanographic conditions in order to maximize the spatial and temporal coexistence of larvae and prey (Dickey-Collas et al. 1996).

Most of the nutrient-rich freshwater discharged from the Mississippi River flows west along the shelf to form a lower salinity coastal boundary layer that extends 10– 20 km from shore (Ichiye 1962; Cochrane and Kelly 1986; Wiseman et al. 1986). Over 50% of this nutrientrich water reaches Texas (Dinnel and Wiseman 1986)

with about a 1-month lag-time (Temple et al. 1977). During late spring/early summer, southeast winds favor weak upwelling of cooler waters off Texas (Fig. 7), which triggers a reversal of coastal boundary layer flow and convergence of upwelling and riverine waters over the broad Louisiana/Texas continental shelf. As a result, lower-salinity nutrient-rich coastal waters turn offshore (Cochrane and Kelly 1986; Dagg et al. 1991), which can enhance the physical stratification of the water column and alter trophic processes [e.g., aggregation of food at intermediate trophic levels (Dagg 1988)]. Concentrations of *Chloroscombrus chrysurus* larvae over the Louisiana/ upper Texas shelf during summer (Fig. 2a) and concentrations of T. lathami near the Mississippi River delta during winter and early spring (Shaw and Drullinger 1990) coincide both spatially and temporally with areas of higher ZDVs (Fig. 6) and regional oceanographic processes that could enhance feeding opportunities for larvae. Similarly, concentrations of Caranx crysos larvae during summer at stations in a hydrographically dynamic area along the outer edge of the continental shelf (Figs. 2c, 6b) are consistent with a linkage between regional oceanographic processes (e.g., shelf mixing of riverine and coastal boundary layer waters) and biological productivity.

Based on the spatial and temporal distributions of carangid larvae in this study, the abundant but unidentified carangid near the Mississippi River delta during early fall (Grimes and Finucane 1991) may be the larvae of *C.crysos*. Similarly, concentrations of an unidentified carangid along the eastern edge of the Loop Current during spring (Richards et al. 1993) may be *D. punctatus*larvae. Concentrations of *D. punctatus* in areas where ZDVs are higher (Farris 1961; Conand and Franqueville 1973; Ortner et al. 1989; Katsuragawa and Matsuura 1992) should not be surprising given that *Decapterus*, *Trachurus*, and *Selar* are planktivorous throughout life.

The small number of larval *C. crysos* taken in SEA-MAP and other Gulf ichthyoplankton surveys (Flores-Coto and Sanchez-Ramirez 1989; Katsuragawa and Matsuura 1992) is surprising given the abundance of adults in the survey area (Klima 1971). Similarly, the

fact that fishery surveys show juvenile and adult *T. lathami,E. bipinnulata, C.hippos/latus, Seriola,* and *Trachinotus* to be more abundant than ichthyoplankton surveys suggest inadequate sampling at appropriate times and locations, gear avoidance, or gear bias. Patterns of abundance vary widely with gear, mesh size, and depth sampled. In theory, oblique bongo-net tows, such as those reported here, sample the water column in an unbiased way (Kingsford 1988). Oblique tows, however, poorly represent species that primarily occupy surface waters because bongo nets under-sample narrow vertical strata. Similarly, larvae of some species may be concentrating in deeper waters or near the bottom beyond depths sampled by SEAMAP protocol.

In conclusion, co-occurrence of concentrations of larvae of some species of carangids with areas of higher ZDVs suggests that these areas serve as important nurseries. While higher total ZDVs are not a direct measure of food availability of the appropriate size and species composition (Sameoto 1984), the co-occurrence of higher concentrations of larvae and zooplankton adds to the probability of successful feeding, growth, and survival (Kingsford 1988; Grimes and Finucane 1991).

Acknowledgements The Marine Fisheries Initiative (MARFIN) Program (contracts NA86-WC-H-06117, NA87WC-H-06135, NA88WC-H-MF198, and NA90AA-H-MF111) funded final analyses. Special thanks to David L. Drullinger and Talat Farooqi at Louisiana State University who helped identify carangid larvae. Thanks also to the SEAMAP Ichthyoplankton Program; to data manager Ken Savastano [NOAA Fisheries, Pascagoula (retired)] for supplying SEAMAP environmental data; and to the Florida Department of Natural Resources for providing access to the SEAMAP samples. Thanks to Phil Caldwell (NOAA Fisheries, Galveston) for plotting the carangid and zooplankton distribution maps.

References

- Aprieto VL (1974) Early development of five carangid fishes of the Gulf of Mexico and the South Atlantic Coast of the United States. Fish Bull US 72: 414–443
- Arnold EL Jr (1958) Gulf of Mexico plankton investigations: 1951–53. Special Scientific Report—Fisheries No. 269. US Fish and Wildlife Service, Washington, D.C.
- Austin HM, Jones JI (1974) Seasonal variation of physical oceanographic parameters on the Florida middle ground and their relation to zooplankton biomass on the West Florida shelf. Fla Sci 37:16–32
- Berry FH (1959) Young jack crevalles (*Caranx* species) off the southeastern Atlantic Coast of the United States. Fish Bull US 152:417–535
- Bogdanov DV, Sokolov VA, Khromov NS (1968) Regions of high biological and commercial productivity in the Gulf of Mexico and Caribbean Sea. Oceanology 8:371–381
- Bullis HR Jr, Carpenter JS (1968) Latent fishery resources of the central West Atlantic region. Univ Wash Publ Fish New Ser 4:61-64
- Bullis HR Jr, Thompson JR (1970) Bureau of commercial fisheries exploratory fishing and gear research base, Pascagoula, Mississippi—July 1, 1967 to June 30, 1969. Circular No. 351, US Fish and Wildlife Service
- Cochrane JD, Kelly FJ (1986) Low-frequency circulation on the Texas-Louisiana continental shelf. J Geophys Res 91:10645– 10659

- Conand F, Franqueville C (1973) Identification and seasonal distribution of carangid larvae off the coasts of Senegal and Gambia (in French). Bull Inst Fondam Afr Noire Ser A Sci Nat XXXV:951–978
- Dagg MJ (1988) Physical and biological responses to the passage of a winter storm in the coastal and inner shelf waters of the northern Gulf of Mexico. Cont Shelf Res 8:167–178
- Dagg MJ, Ortner PB, Al-Yamani FY (1987) Winter-time distribution and abundance of copepod nauplii in the northern Gulf of Mexico. Fish Bull US 86:319–330
- Dagg M, Grimes C, Lohrenz S, McKee B, Twilley R, Wiseman W Jr (1991) Continental shelf food chains of the northern Gulf of Mexico. In: Sherman K, Alexander LM, Gold BD (eds) Food chains, yields, models, and management of large marine ecosystems. Westview, Boulder, Colo., pp 67–106
- Daniel WW (1990) Applied nonparametric statistics, 2nd edn. PWS-Kent, Boston, Mass.
- Dickey-Collas MR, Gowen RJ, Fox CJ (1996) Distribution of larval and juvenile fish in the western Irish Sea: relationship to phytoplankton, zooplankton biomass and recurrent physical features. Mar Freshw Res 47:169–181
- Dinnel SP, Wiseman WJ Jr (1986) Fresh water on the Louisiana and Texas shelf. Cont Shelf Res 6:765–784
- Ditty JG, Truesdale FM (1984) Ichthyoplankton surveys of near shore Gulf waters off Louisiana: January–February and July, 1976. Assoc SE Biol Bull 31:55–56
- Ditty JG, Zieske GG, Shaw RF (1988) Seasonality, depth, and areal distribution of larval fishes in the northern Gulf of Mexico above 26° 00′ N latitude. Fish Bull 86:811–823
- Elwertowski J, Boely T (1971) Repartition saisonniere' des poissons pelagiques cotiers dans les eaux Mauritaniennes et Senegalaises (in French). Trav Doc ORSTOM, Dakar-Thiaroye, No. 22
- Fahay MP (1975) An annotated list of larval and juvenile fishes captured with surface-towed meter net in the South Atlantic Bight during four RV Dolphin cruises between May 1967 and February 1968. NOAA Tech Rep NMFS SSRF-685
- Farris D (1961) Abundance and distribution of eggs and larvae and survival of larvae of jack mackerel (*Trachurus symmetricus*). Fish Bull US 61:247–279
- Fields HM (1962) Pompanos (*Trachinotus* spp.) of South Atlantic Coast of the United States. Fish Bull US 207:189–222
- Finucane JH, Collins LA, Barger LE (1979a) Determine the effects of discharges on seasonal abundance, distribution, and composition of ichthyoplankton in the oil field. In: Jackson WB (ed) Environmental assessment of an active oil field in the northwestern Gulf of Mexico, 1977–1978, vol 2. Data management and biological investigation. NOAA Final Report to EPA, Contract No EPA-IAG-D5-E693-EO, pp 2.3.6.-1 to 2.3.6.-157
- Finucane JH, Collins LA, Barger LE, McEachran JB (1979b) Ichthyoplankton/mackerel eggs and larvae. In: Jackson WB (ed) Environmental studies of the south Texas outer continental shelf 1977. NOAA Final Report to BLM, Contract No AA550-IA7-21, pp 1-504
- Flores-Coto C, Sanchez-Ramirez M (1989) Larval distribution and abundance of Carangidae (Pisces) from the southern Gulf of Mexico. Gulf Res Rep 8:117–128
- Flores-Coto C, Sanchez-Rameriz M, Zavala-Garcia F (1998) Desarrollo larvario de *Hemicaranx amblyrhynchus* (Pisces: Carangidae), del sur del Golfo de Mexico. Rev Biol Trop 46:431–438
- Flores-Coto C, Flores-Hernandez F, Zavala-Garcia F, Arenas-Fuentes V, Monreal-Gomez MA, Salas-de-Leon DA (2000) Oceanic and neritic ichthyoplankton at the edge of the continental shelf in the southern Gulf of Mexico. Gulf Carib Res 12:31–35
- Flores-Coto C, Rivas-Vega R, Zavala-Garcia F, Sanchez-Robles J (2001) Vertical distribution of larval carangids in the southern Gulf of Mexico. Gulf Carib Res 13:1-8
- Goodwin JM, Finucane JH (1985) Reproductive biology of blue runner (*Caranx crysos*) from the eastern Gulf of Mexico. Northeast Gulf Sci 7:139–146
- Grimes CB, Finucane JH (1991) Spatial distribution and abundance of larval and juvenile fish, chlorophyll and macrozoo-

- plankton around the Mississippi River discharge plume, and the role of the plume in fish recruitment. Mar Ecol Prog Ser 75:109–119
- Hales LS Jr (1987) Distribution, abundance, reproduction, food habits, age, and growth of round scad, *Decapterus punctatus*, in the South Atlantic Bight. Fish Bull US 85:251–268
- Hoese HD (1965) Spawing of marine fishes in Port Aransas, Texas, as determined by the distribution of young, and larvae. PhD dissertation, University of Texas
- Houde ED (1982) Kinds, distributions and abundances of sea bass larvae (Pisces: Serranidae) from the eastern Gulf of Mexico. Bull Mar Sci 32:511–522
- Houde ED, Chitty N (1976) Seasonal abundance and distribution of zooplankton, fish eggs, and fish larvae in the eastern Gulf of Mexico, 1972–74. NOAA Tech Rep NMFS SSRF-701
- Houde ED, Leak JA, Dowd CE, Berkeley SA, Richards WJ (1979) Ichthyoplankton abundance and diversity in the eastern Gulf of Mexico. University of Miami, Coral Gables, Fla.
- Howey TW (1976) Plankton of the Gulf of Mexico: distribution of displacement volume, occurrence of systematic groups, abundance and diversity among copepods. PhD dissertation, Department of Zoology, Louisiana State University, Baton Rouge
- Ichiye T (1962) Circulation and water mass distribution in the Gulf of Mexico. Geofis Int (Mexico City) 2:47–76
- Juhl R (1966) Experimental fish trawling survey along the Florida west coast. Commer Fish Rev 28:1–5
- Katsuragawa M, Matsuura Y (1992) Distribution and abundance of carangid larvae in the southeastern Brazilian Bight during 1975–1981. Bolm Inst Oceanogr Sao Paulo 40:55–78
- Khromov NS (1969) Distribution of plankton in the Gulf of Mexico and some aspects of its seasonal dynamics (English translation). In: Bogdanov AS (ed) Soviet-Cuban fishery research. Israel Program for ScientificTranslations, Jerusalem, pp 36–56
- Kingsford MJ (1988) The early life history of fish in coastal waters of northern New Zealand: a review. NZ J Mar Freshw Res 22:463–479
- Klima EF (1971) Distribution of some coastal pelagic fishes in the western Atlantic. Commer Fish Rev 33:21–34
- Laroche WA, Smith-Vaniz WF, Richardson SL (1984) Carangidae:
 development. In: Moser HG, Richards WJ, Cohen DM, Fahay
 MP, Kendall AW Jr, Richardson SL (eds) Ontogeny and Systematics of Fishes. American Society of Ichthyologists and Herpetologists Special Publication No. 1. Allen, Lawrence, Kan., pp 510–522
- Leak JC (1977) Distribution and abundance of Carangidae (Pisces, Perciformes) larvae in the eastern Gulf of Mexico, 1971–1974. MSc thesis, University of Miami, Coral Gables, Fla.
- Leak JC (1981) Distribution and abundance of carangid fish larvae in the eastern Gulf of Mexico, 1971–1974. Biol Oceanogr 1:1–28
- Leffler DL, Shaw RF (1992) Age validation, growth, and mortality of larval Atlantic bumper (Carangidae: *Chloroscombrus chrysurus*) in the northern Gulf of Mexico. Fish Bull US 90:711–719
- Lyons JM (1978) Distribution and abundance of the larvae of Decapterus punctatus (Pisces, Carangidae) and Bothus spp. (Pisces, Bothidae) in the Florida and Yucatan Straits. MSc thesis, Florida State University, Tallahassee
- Masuma S, Kanematu M, Teruya K (1990) Embryonic and morphological development of larvae and juveniles of the amberjack, *Seriola dumerili*. Jap J Ichthyol 37:164–169
- McBride RS, Stengard FJ, Mahmoudi B (2002) Maturation and diel reproductive periodicity of round scad (Carangidae: *Decapterus punctatus*). Mar Biol 140:713–722
- McKenney TW, Alexander EC, Voss GL (1958) Early development and larval distribution of the carangid fish, *Caranx crysos* (Mitchill). Bull Mar Sci 8:167–199

- Montolio MA (1976) Estudio taxonomico y morfometrico de los estadios larvales de dos especies de Carangidae *Decapterus* punctatus (Agassiz, 1829) y Caranx crysos (Mitchill, 1815) y su distribucion en el Golfo de Mexico (in Spanish). Rev Invest Inst Nac Pesca 2:85–125
- Nakamura EL (1980) Carangids of the northern Gulf of Mexico. In: Flandorfer M, Skupien L (eds) Proceedings of a workshop for potential fishery resources of the northern Gulf of Mexico, March 4–5, 1980, New Orleans, La., pp 18–33
- Ortner PB, Hill LC, Cummings SR (1989) Zooplankton community structure and copepod species composition in the northern Gulf of Mexico. Cont Shelf Res 9:387–402
- Reintjes JW (1979) Coastal herrings and associated species: a profile of species or groups of species, their biology, ecology, current exploitation with economic and social information. Report to the Gulf of Mexico Fish Management Council by NOAA NMFS Southeast Fish Science Center, Beaufort, North Carolina
- Richards WJ, McGowan MF (1989) Biological productivity in the Gulf of Mexico: identifying the causes of variability in fisheries. In: Sherman K, Alexander LM (eds) Biomass yields and geography of large marine ecosystems. American Association for the Advancement of Science, Symposium No. 111, pp 287–325
- Richards WJ, McGowan MF, Leming T, Lamkin JT, Kelley S (1993) Larval fish assemblages at the Loop Current boundary in the Gulf of Mexico. Bull Mar Sci 53:475–537
- Ruple DL (1984) Occurrence of larval fishes in the surf zone of a northern Gulf of Mexico barrier island. Estuar Coast Shelf Sci 18:191–208
- Sameoto DD (1984) Environmental factors influencing diurnal distribution of zooplankton and ichthyoplankton. J Plankton Res 6:767–791
- Sanchez-Ramirez M, C. Flores-Coto (1993) Desarrollo larvario y clave de identificación de algunas espesies de la familia Carangidae (Pisces) del sur del Golfo de Mexico. An Inst Cienc Mar Limnol Univ Nac Auton Mex 20:1–24
- Sanchez-Ramirez M, Flores-Coto C (1998) Growth and mortality of larval Atlantic bumper *Chloroscombrus chrysurus* (Pisces: Carangidae) in the southern Gulf of Mexico. Bull Mar Sci 63:295–303
- Sanchez-Velasco L, Flores-Coto C (1994) Larval fish assemblages at the Yucatan Shelf and in the Mexican Caribbean Sea during the upwelling period (Spring, 1985). Sci Mar 58:289–297
- Shaw RF, Drullinger DL (1990) Early-life-history profiles, seasonal abundance, and distribution of four species of carangid larvae off Louisiana, 1982 and 1983. NOAA NMFS Technical Report No. 89
- Smith PE, Richardson SL (1977) Standard techniques for pelagic fish egg and larva surveys. FAO Fisheries Technical Paper No. 175
- Tachihara K, Ebisu R, Tukashima Y (1993) Spawning, eggs, larvae and juveniles of the purplish amberjack, Seriola dumerili. Nippon Suisan Gakkaishi 59:1479–1488
- Temple RF, Harrington DL, Martin JA (1977) Monthly temperature and salinity measurements of continental shelf waters in the western Gulf of Mexico 1963–1965. NOAA Technical Report NMFS SSRF-707
- Thompson BA, Brown L (1994) Identification, distribution and life history of lesser amberjack, *Seriola fasciata* (abstract). Annual Meeting of the American Fisheries Society, Louisiana and Mississippi Chapters, February 10–11, Natchez, Miss.
- Thompson BA, Beasley M, Wilson CA (1998) Age distribution and growth of greater amberjack, *Seriola dumerili*, from the north-central Gulf of Mexico. Fish Bull US 97:362–371
- Wiseman WJ Jr, Turner RE, Kelly FJ, Rouse LJ Jr, Shaw RF (1986) Analysis of biological and chemical associations near a turbid coastal front during winter 1982. Contrib Mar Sci, Univ Texas 29:141–151